

Recovery of plant diversity and soil nutrients during stand development in subtropical forests of Mizoram, Northeast India

SH. B. SINGH¹, B.P. MISHRA², S.K. TRIPATHI^{1,✉}

¹Department of Forestry, Mizoram University, Aizawl-796004, Mizoram, India. Tel. +91-389-2330394, Fax. +91-389-2330834, ✉email: sk_tripathi@rediffmail.com

²Department of Environmental Science, Mizoram University, Aizawl-796004, Mizoram, India

Manuscript received: 4 January 2015. Revision accepted: 28 August 2015.

Abstract. Singh ShB, Mishra BP, Tripathi SK. 2015. Recovery of plant diversity and soil nutrients during stand development in subtropical forests of Mizoram, Northeast India. *Biodiversitas* 16: 205-212. The present study assessed the recovery of tree species diversity and soil nutrient dynamics with stand development in subtropical semi-evergreen forest of Mizoram. The study was carried out in two regenerating forest stands following disturbance and one undisturbed forest. *Schima wallichii* was the dominant species in all stands showing IVI of 63.8, 83.3 and 75.9 in undisturbed, moderately disturbed and highly disturbed stands, respectively. *Castanopsis tribuloides* was co-dominant species in the undisturbed and the moderately disturbed but this species was replaced by *Sterculia villosa* in the highly disturbed stand. The shift in position of species and families from undisturbed to highly disturbed stands could be linked with degree of disturbance. Log-normal dominance-distribution curves in the undisturbed and moderately disturbed stand indicating the stability of community, while short hooked curve in the highly disturbed stand indicates unstable nature of the community. The soil properties (organic C, total nitrogen, and available phosphorus) increased significantly during the course of stand development, whereas, decrease with the depth in these forest stands.

Keywords: Community, disturbance, plant diversity, tropical semi-evergreen forest

INTRODUCTION

Biodiversity provides many essential goods and services to the mankind particularly in densely populated tropical ecosystems. Rapid decrease in tropical biodiversity is recorded in many forest ecosystems over the world as a result of human activities like habitat destruction, over exploitation, pollution and species introduction (Pimm et al. 1995; Pragasan and Parthasarthy 2010). These human activities have led to the conversion of natural forest areas into different stages of forest degradation that exhibited significantly different structural and functional characteristics at local and regional levels (Sapkota et al. 2010). Therefore, proper species diversity along with other characteristics need to be recorded in different ecosystems to account multi-dimensional aspect of biodiversity that should take into account the variation in environmental factors, taxonomic variations among species, compositional diversity and functional diversity (Roy et al. 2004). The extents of deforestation in recent years in tropical regions draw attention to the urgent need for intervention to restore and protect biodiversity, ecological functioning and the supply of goods and services used by poor rural communities (Lamb et al. 2005; Cayuela et al. 2006).

The North-east India is known for its rich biodiversity with high level of endemism (Khan et al. 1997). Several studies have been carried out to quantify plant diversity and to understand the ecology of forest communities of

different areas (Mishra et al. 2004; Laloo et al. 2006; Mishra 2010; Tynsong and Tiwari 2010, 2011). The plant inventory and community characteristics of tropical semi-evergreen forest of Mizoram have been studied by some workers (Lalramnghinglova 2003; Singh et al. 2011, 2012). In Mizoram, one of the states of North-east India, an extensive sandstone quarry has been carried out for the many developmental activities like construction of roads and buildings in the last few decades that converted large tract of forest area into degraded forest and created an unfavorable habitat conditions for plant growth and development. The unfavorable habitat conditions prevailing in the mined areas have reduced the chances of regeneration of many species, thereby reducing the number of species in mined areas.

Anthropogenic disturbance is one of the significant impacts arising out of mining and quarrying activities and is mainly in the form of alteration of land structure due to excavation, stacking of top soil and loss of fertile soil due to dumping of mined overburden on top soil. Sandstone mining causes damage to property, depletion of forest land, adverse effects on the aquatic biodiversity and public health. The present study aimed to determine the recovery of plant community characteristics and soil nutrients in secondary successional forests with stand development in relation to a reference forest in semi-evergreen tropical region of Mizoram, Northeast India.

MATERIAL AND METHODS

The study was conducted during 2009 to 2011 in the forest patches within and outside the Mizoram University campus situated in Tanhril area (23°45'25" N-23°43'37" N latitude and 92°38'39" E-92°40'23" E longitude) of Aizawl District, Mizoram, India (Figure 1). For detailed investigation, a total of three study sites were selected along age gradient after disturbance in terms of sandstone quarry, representing undisturbed (UD), moderately disturbed (MD) and highly disturbed (HD) stands. The natural forest stand where no sandstone mining activity was done in the past referred to as undisturbed stand. The moderately disturbed forest stand is a secondary forest developed naturally after the abandonment of sandstone quarry (mining operation about 7 years back in 2002). The highly disturbed forest stand is an open mining area where sandstone mining is continued. Tree species (15 cm girth at breast height, GBH, i.e. 1.3 m above ground) were recorded using 70 quadrats of 10 x 10m size at each site. The plant species were identified with the help of herbarium of the concerned University Department; herbarium of the Botanical Survey of India (BSI), Eastern circle, Shillong, and counter checked with the help of regional floras (Kanjilal et al. 1934-1940; Haridasan and Rao 1985). The field data on vegetation was quantitatively analyzed for phyto-sociological attributes namely, frequency, density and abundance as proposed by Curtis and McIntosh (1950). The Importance value index (IVI) was determined as per Philips (1959). Species diversity and dominance indices were determined following the methods as outlined in Misra (1968); Mueller-Dombois and

Ellenberg (1974). The soil samples were collected with the help of soil corer for top-soil (0-10 cm depth) and sub-soil (10-20 cm depth). The physico-chemical characteristics of soil have been analyzed by the methods outlined by Allen et al. (1976) and Anderson and Ingram (1993). Correlation was also developed between different vegetational and soil parameters across different sites.

RESULTS AND DISCUSSION

Plant diversity and community characteristics

Altogether, a total of 71 woody plant species belonging to 59 genera and 33 families of angiosperms were recorded from undisturbed, moderately disturbed and highly disturbed forest stands. Of this, 50 species representing 43 genera and 29 families, 49 species belonging to 41 genera and 22 families and 24 species belonging to 21 genera and 17 families were reported from the undisturbed, moderately disturbed and highly disturbed stand, respectively. It was found that *Wendlandia tinctoria* (Roxb.) DC., *Schima wallichii* (DC.) Korthals., *Embllica officinalis* Gaertn., *Callicarpa arborea* Roxb., *Castanopsis tribuloides* DC., and *Sterculia villosa* Roxb. ex Smith species were common in all three stands, and *Acacia farnesiana*, *Debregeasia velutina*, *Murraya koenigii*, *Turpinia nepalensis* and species *Cassia laevigata*, *Melia azedarach*, *Pterospermum acerifolium* and species *Lepionurus oblongifolius*, *Ficus virens*, *Grewia macrophylla* were restricted to undisturbed, moderately disturbed and highly disturbed stand, respectively. There was shift in dominance of plant species and families from undisturbed to highly disturbed




Figure 1. Map of Mizoram, India showing the study area

stand. *S. wallichii* was the dominant species (IVI 63.76) in the undisturbed stand. The co-dominant species were *C. tribuloides* (IVI 19.99) and *C. arborea* (IVI 19.44). In the moderately disturbed stand, *S. wallichii* was dominant species (IVI 83.31) and it was followed by co-dominant species namely, *C. tribuloides* (IVI 39.81) and *C. arborea* (IVI 17.54). In the highly disturbed stand, *S. wallichii* was recorded as a dominant species (IVI 75.87), and it was followed by co-dominant species namely, *S. villosa* (IVI 39.04) and *C. arborea* (IVI 24.89). *S. wallichii*, the dominant species in all three stands, contributed maximum tree density in the undisturbed stand (225 ind. ha⁻¹ and basal area 5.04 m²ha⁻¹), moderately disturbed stand (222 ind. ha⁻¹ and basal area 3.28 m²ha⁻¹) and highly disturbed stand (105 ind. ha⁻¹ and basal area 1.59 m²ha⁻¹) (Table 1). The distribution pattern of species differed from undisturbed to the highly disturbed stand. There was preponderance of contagious distribution. Whereas, few species namely, *C. arborea* and *E. officinalis* showed random distribution.

Euphorbiaceae was the dominant family with maximum number of species in the undisturbed (5) and moderately disturbed (5) stands. However, it was replaced by Mimosaceae (3) in the highly disturbed stand. The co-dominant families were Fagaceae (4), Lauraceae (4), Mimosaceae (4) in the undisturbed stand; Caesalpiniaceae (4), Fagaceae (4), Lauraceae (4) in the moderately disturbed stand; Anacardiaceae (2), Rubiaceae (2), Papilionaceae (2) in the highly disturbed stand (Table 2).

There was sharp decline in tree density from the undisturbed stand (970±3.2 ind. ha⁻¹) to the moderately disturbed stand (742±3.4 ind. ha⁻¹) and finally to the highly disturbed stand (410±3.2 ind. ha⁻¹). A similar trend in results was also observed for total tree basal area which was highest in the undisturbed stand (20.8 m² ha⁻¹) followed by moderately disturbed stand (9.85 m² ha⁻¹) and highly disturbed stand (5.38 m² ha⁻¹). Shannon-Wiener diversity index was maximum (3.3) in the undisturbed stand and minimum in the highly disturbed stand (2.6). The species richness Index was highest in moderately disturbed stand followed by undisturbed stand and highly disturbed stands. The Evenness Index did not show much variation in results with respect to the degree of disturbance. The Sorenson's Index of similarity for tree species was maximum (0.58) between undisturbed and moderately disturbed stands. On contrary, value was minimum (0.46) between moderately disturbed and highly disturbed stands (Table 3).

Findings on the girth class distribution of trees showed a decreasing trend in the number of individuals from lower to higher girth classes, irrespective of stands, except in case of the undisturbed stand where number of individuals was highest in girth classes 50-70 cm and 30-50 cm. The moderately disturbed stand showed maximum number of adult trees (girth classes 30-50 cm). However, mature trees (girth classes 50-70 cm) were maximum in the undisturbed stand. There was a sharp decline in the number of individuals in the higher girth classes with increase in degree of disturbance (Figure 2).

The log-normal dominance-diversity curve (based on IVI) was found in the undisturbed and moderately disturbed stands; however, it was short hooked in case of highly disturbed stand (Figure 3).

Soil characteristics

The average soil moisture on different stands ranged from 23.66 to 28.84%. pH ranged from 4.45 to 6.18 which indicated that the soils on all stands were acidic. Compared to the moderately disturbed and highly disturbed stands, the undisturbed stand has the highest values of soil in organic carbon, total nitrogen, available phosphorus, exchangeable potassium as given in Table 4.

Correlation between vegetation and soil parameters

The correlation between different vegetation parameters along disturbance gradient is given in Figure 4. The tree density was positively correlated with the tree basal area, tree diversity and tree richness (Figure 4 A, B, D), whereas, tree basal area was negatively correlated with concentration of dominance (Figure 4). Tree richness was positively correlated with diversity and negatively with concentration of dominance but the relationship was rather weak (Figure 4 E, F).




Figure 2. Distribution of tree species in different girth classes in the undisturbed (UD), moderately disturbed (MD) and highly disturbed (HD) forest stands




Figure 3. Dominance-diversity curves of tree species along disturbance gradient




Figure 4. Correlation between various vegetational parameters along disturbance gradient




Figure 5. Correlation between various soil parameters for top-soil along disturbance gradient

The correlation between various soil parameters for top-soil and sub-soil along disturbance gradient is given in Figure 5 and 6. The soil moisture content showed a positive correlation with soil organic carbon and negative correlation with pH (Figure 5 A, B). However, soil organic carbon showed significant and positive correlation with total nitrogen and exchangeable potassium (Figure 5 C, E).

Available phosphorus showed positive and significant correlation with exchangeable potassium, and the total nitrogen showed positive correlation with available phosphorus (Figure 5 D, F). Correlation among different nutrients showed similar trends in top-and sub-soil with varying degree of coefficients (Figure 5, 6).




Figure 6. Correlation between various soil parameters for sub-soil along disturbance gradient

Discussion

The sandstone mining impacted vegetation composition, community organization to a great extent, and there was decrease in species richness, tree density and basal area from undisturbed to the highly disturbed stand. *S. wallichii* was the dominant tree species in all the stands. However, *C. tribuloides*, the co-dominant species of the undisturbed and moderately disturbed stands was replaced by *S. villosa* in the highly disturbed stand. Moreover, the species tolerant to stress showing better growth and survival under disturbed condition, such species express greater IVI in the disturbed stand. Similarly, Euphorbiaceae was the dominant family in both undisturbed and moderately disturbed stands, whereas, it was replaced by Mimosaceae that was the dominant family in the highly disturbed stand. The shift in position of the species and families along the disturbance gradient could be linked with the levels of anthropogenic disturbance and similar trends was also reported by number of workers in the past (Kadavul and Parthasarathy 1999; Mishra et al. 2003, 2004, 2005; Mishra and Laloo 2006).

During present investigation, it has been found that majority of the species showed contagious distribution pattern and few were distributed randomly. Similar, distribution pattern has also been reported by other workers (Singh and Yadav 1974; Metha et al. 1997). The trend in tree population structure observed during present study is similar to those reported from the forest at Costa Rica (Nadakarni et al. 1995), Brazilian Amazon (Campbell et al. 1992), Eastern Ghats (Kadavul and Parthasarathy 1999) and sub-tropical humid forest of Meghalaya (Mishra et al. 2004). The log-normal dominance distribution curve in the

undisturbed and moderately disturbed stands indicates stable community. However, short hooked curve in the highly disturbed stand depicts un-stable community. The past workers have also reported a similar trend in results (Khera et al. 2001; Tripathi et al. 2004; Mishra et al. 2004, 2005).

The soil properties greatly varied from undisturbed to highly disturbed stand. High pH value in highly disturbed stand could be due to low accumulation of decomposed organic matter and extraction of sand stone. The soil organic carbon, total nitrogen, available phosphorus and exchangeable potassium varied greatly along disturbance gradient and decreased from undisturbed to highly disturbed stand and higher values were reported from top-soil. This could be linked with presence of thick humus layer on forest floor of the undisturbed stand. Litter accumulation on forest floor is positively linked with litter decomposition and plays a significant role in maintenance of soil moisture content and other micro-environmental conditions (Arunachalam and Pandey 2003; Reddy 2010; Mishra 2010; Tripathi et al. 2012). Nayak and Srivastava (1995) have also reported a similar trend in results from the humid sub-tropical soils in north east India.

A positive correlation of species richness with diversity index was observed with few exceptions, and findings are in conformity with the report of Tripathi et al. (1989) that the species richness is positively correlated with Shannon diversity index in majority of cases in western Himalayan forests. Various diversity indices also followed similar trend (Jha et al. 2005; Yu-Hai et al. 2009). A positive correlation was observed between total basal area and organic carbon in top-soil and sub-soil. The findings are supported by Sharma et al. (2009).

Table 1. Phyto-sociological attributes of tree species along disturbance gradient

Species	Undisturbed		Moderately disturbed		Highly disturbed	
	Density (Ind. ha ⁻¹)	IVI	Density (Ind. ha ⁻¹)	IVI	Density (Ind. ha ⁻¹)	IVI
<i>Acacia auriculiformis</i> A. Cunn. ex Benth.	-	-	3	1.36		
<i>Acacia farnesiana</i> (L.) Willd.	5	1.17	-	-		
<i>Albizia chinensis</i> Osb. Merr.			4	2.04	15	10.19
<i>Albizia lebbbeck</i> (L.) Benth.	8	2.37	13	4.95	6	4.78
<i>Albizia procera</i> (R.) Benth.	10	3.73	12	6.39	12	9.38
<i>Antidesma diandrum</i> B.Hey. ex (Roxb.)	18	6.39	2	0.96		
<i>Aporosa roxburghii</i> Baill	3	0.88				
<i>Artocarpus heterophyllus</i> Lam	4	1.53				
<i>Bauhinia purpurea</i> L.	3	1.21				
<i>Beilschmiedia assamica</i> Meisn.	7	1.53				
<i>Bombax ceiba</i> L.	7	2.24			4	3.48
<i>Bridelia cuneata</i> Gehrms	15	4.99				
<i>Bridelia stipularis</i> Bl.	20	6.3	6	2.56		
<i>Bursera serrata</i> Wall.ex Colebr.	2	0.78				
<i>Callicarpa arborea</i> Roxb.	58	19.44	42	17.54	32	24.89
<i>Callicarpa macrophylla</i> Vahl.	4	1.32				
<i>Cassia laevigata</i> Willd			5	1.73		
<i>Castanopsis hystrix</i> A.DC.	31	10.41	8	3.2		
<i>Castanopsis tribuloides</i> A.DC.	62	19.99	102	39.81	24	18.05
<i>Clausena excavata</i> Burn.f.			5	2.95		
<i>Colona floribunda</i> (Wall.ex Kurz) Craib					7	6.23
<i>Combretum dasystachyum</i> Kurz.	18	4.92				
<i>Crotalaria linifolia</i> L.	13	3.5	4	2.45		
<i>Debregeasia velutina</i> Gaud.	11	4.96				
<i>Delonix regia</i> (Boj)Rafin.			1	0.86		
<i>Derris robusta</i> Roxb.exDC.			2	0.93	6	4.44
<i>Dipterocarpus turbinatus</i> Gaertn.	7	2.26			1	1.77
<i>Emblica officinalis</i> Gaertn.	32	10.45	30	11.87	20	13.78
<i>Engelhardia spicata</i> Lechen ex Bl.	2	0.72				
<i>Erythrina stricta</i> Roxb.	15	5.17			4	2.96
<i>Eucalyptus globulus</i> Labill.			1	0.75		
<i>Eugenia macrocarpa</i> Schlttdl. & Cham.	20	6.05				
<i>Eugenia malaccensis</i> L.			5	2.18		
<i>Ficus elastica</i> Roxb.ex Hornem			6	2.83		
<i>Ficus virens</i> Ait.	6	2.45			5	4.83
<i>Glochidion velutinum</i> Wt.	5	1.53	15	6.14	1	1.77
<i>Grevillea robusta</i> A.Cunn.ex R.Br.			2	0.75		
<i>Grewia macrophylla</i> G.Don	2	0.54			5	3.22
<i>Lagerstroemia parviflora</i> Roxb.	2	1.17	2	0.91	5	4.19
<i>Lepionurus oblongifolius</i> (Griff.) Mast	8	2.42			8	5.09
<i>Litsea chinensis</i> (Lour.) C.B.Rob.	6	1.75	14	5.83		
<i>Litsea citrata</i> Bl.	15	4.73	8	3.63		
<i>Macaranga denticulata</i> (Bl.) Mull. Arg.			2	1.06		
<i>Machilus bombycina</i> King ex Hook.f.			4	1.57		
<i>Mangifera indica</i> Linn.			4	2.29		
<i>Melia azedarach</i> Linn.			5	2.07		
<i>Meliosma pinnata</i> Roxb.	18	5.04	5	2.09		
<i>Michelia champaca</i> Linn.			3	1.36		
<i>Michelia panduana</i> Hook.Thamn.			5	1.95		
<i>Micromelum integerrimum</i> (B-Ht.ex Can.)Wt.	11	3.8	5	1.78		
<i>Murraya koenigii</i> (L.)Spreng.	15	3.35	17	7.11		
<i>Oroxylum indicum</i> Benth.ex Kurz.	9	2.33				
<i>Phoebe cooperiana</i> Pc. Kanj. & Das.	14	4.98	8	4.09		
<i>Plumeria acutifolia</i> Poir.			2	1.42		
<i>Psidium guajava</i> Linn.	4	1.48	5	3.08		
<i>Pterospermum acerifolium</i> Linn.	9	3.38				
<i>Quercus semiserrata</i> Roxb.	14	4.63	10	3.84	7	5.04
<i>Quercus spicata</i> Sm.	32	8.72	2	0.86		
<i>Rhus succedanea</i> Linn.	21	7.44	22	8.11	21	14.67
<i>Sarcochlamys pulcherrima</i> Roxb.			2	1.29		
<i>Saurauia napaulensis</i> D.C.			12	5.07		
<i>Schefflera wallichiana</i> (Wight & Arn.) Harms	18	5.76	2	1.5		
<i>Schima wallichii</i> (DC.) Korthals.	225	63.76	222	83.31	105	75.87
<i>Sterculia villosa</i> Roxb.ex Smith.	57	17.07	24	10.15	58	39.04
<i>Syzygium cumini</i> Linn.			2	0.61		
<i>Tapiria hirsuta</i> Hook.f	17	6.18	2	0.86	9	6.7
<i>Taxus wallichiana</i> Zucc.	12	3.35				
<i>Terminalia arjuna</i> Roxb.W & A	3	1.27				
<i>Toona ciliata</i> M.Roem.	15	5.24	12	6.02	12	10.33
<i>Turpinia nepalensis</i> Wall.ex Wight & Arn.	7	1.83				
<i>Wendlandia paniculata</i> (Roxb.) DC.	26	7.34	28	10.02	26	17
<i>Wendlandia tinctoria</i> (Roxb.) DC.	33	9.53	36	12.49	17	12.3

Table 2. Ranking of families of angiosperm in the undisturbed, moderately disturbed and highly disturbed forest stands

Family rank	Forest stands					
	Undisturbed		Moderately disturbed		Highly disturbed	
	Family	No. of species	Family	No. of species	Family	No. of species
1	Euphorbiaceae	5	Euphorbiaceae	5	Mimosaceae	3
2	Fagaceae	4	Fagaceae	4	Anacardiaceae	2
3	Lauraceae	4	Mimosaceae	4	Fagaceae	2
4	Mimosaceae	3	Lauraceae	4	Papilionaceae	2
5	Caesalpiniaceae	2	Myrtaceae	4	Rubiaceae	2
6	Combretaceae	2	Caesalpiniaceae	4	Tiliaceae	2
7	Moraceae	2	Anacardiaceae	3	Verbenaceae	1
8	Myrtaceae	2	Rubiaceae	3	Dipterocarpaceae	1
9	Verbenaceae	2	Rutaceae	2	Euphorbiaceae	1
10	Anacardiaceae	2	Meliaceae	2	Sterculiaceae	1
11	Papilionaceae	2	Sterculiaceae	2	Caesalpiniaceae	1
12	Bombacaceae	2	Papilionaceae	2	Bombacaceae	1
13	Rubiaceae	2	Theaceae	1	Lythraceae	1
14	Rutaceae	2	Lythraceae	1	Theaceae	1
15	Tiliaceae	1	Magnoliaceae	1	Meliaceae	1
16	Urticaceae	1	Verbenaceae	1	Olacaceae	1
17	Olacaceae	1	Apocynaceae	1	Moraceae	1
18	Lythraceae	1	Araliaceae	1		
19	Juglandaceae	1	Sabiaceae	1		
20	Meliaceae	1	Proteaceae	1		
21	Sapindaceae	1	Saurauaceae	1		
22	Dipterocarpaceae	1	Moraceae	1		
23	Sterculiaceae	1				
24	Taxaceae	1				
25	Theaceae	1				
26	Burseraceae	1				
27	Araliaceae	1				
28	Bignoniaceae	1				
29	Sabiaceae	1				

Table 3. Tree community structure in the undisturbed, moderately disturbed and highly disturbed forest stands

Parameter	Forest stands		
	Undisturbed	Moderately disturbed	Highly disturbed
No. of family	29	22	17
No. of genera	43	41	21
No. of species	50	49	24
Tree density (Indv.ha ⁻¹)	970±3.2	742±3.4	410±3.2
Tree basal area (m ² h ⁻¹)	20.83	9.85	5.38
Shannon-Wiener index	3.3	2.9	2.6
Simpson dominance index	0.07	0.12	0.1
Simpson index of diversity	0.9	0.8	0.8
Simpson reciprocal index	13.3	8.2	9.1
Species richness (Margalef's index)	7.5	7.7	4
Evenness index (Pielou)	0.84	0.74	0.81

Table 4. Physico-chemical characteristics of soil along the disturbance gradient

Stands	Depth (cm)	Moisture content (%)	pH (H ₂ O)	Org-C (%)	TN (%)	Avail P µg g ⁻¹	K µg g ⁻¹
Undisturbed	0-10	Mean	30.45	4.57	2.5	0.26	26.08
		SE ±	2.47	0.08	0.14	0.02	1.36
	10-20	Mean	27.23	4.34	1.89	0.2	17.41
		SE ±	2.65	0.09	0.15	0.02	1.88
Moderately disturbed	0-10	Mean	27.28	5.5	1.8	0.2	14.41
		SE ±	1.56	0.04	0.16	0.02	1.16
	10-20	Mean	24.15	5.1	1.55	0.16	12.16
		SE ±	1.57	0.04	0.12	0.01	1.15
Highly disturbed	0-10	Mean	25.79	6.35	1	0.29	10.25
		SE ±	2	0.04	0.14	0.17	1.43
	10-20	Mean	21.53	6.01	0.76	0.48	6.75
		SE ±	2.28	0.05	0.06	0.23	0.49

Present investigation showed that there is a considerable reduction in the number of species from natural forest to disturbed sites due to unfavorable habitat conditions for plants created by disturbance. The disturbance adversely affected the juvenile stage, leading to arrested survival and growth of seedlings and saplings. The findings of the present study would be an important tool for formulation of appropriate strategies for management of abandoned mined areas through re-vegetation with suitable dominant and co-dominant species of the present study for effective management strategy. Such site-specific selection of species will add a new dimension for rehabilitating of abandoned land and conservation of biodiversity.

ACKNOWLEDGEMENTS

Authors are grateful to the University Grants Commission, New Delhi, India for financial assistant and to the various communities for extending facilities to complete this piece of work desirably.

REFERENCES

- Allen SE, Grimshaw HM, Rowland AP. 1976. Chemical analysis. In: Moore PD, Chapman SB (eds). *Methods in Plant Ecology*. Blackwell, Oxford.
- Anderson JM, Ingram JSI. 1993. *Tropical Soil Biology and Fertility. A Handbook of Methods*. 2nd ed. CAB International, Wallingford, U.K.
- Arunachalam A, Pandey HN. 2003. Ecosystem restoration of jhum fallows in northeast India: microbial C & N along altitudinal and successional gradient. *Restor Ecol* 11: 168-173.
- Campbell DG, Stone JL, Rosas A Jr 1992. A comparison of the phytosociology and dynamics of three floodplain (Varzea) forest of known ages, Rio Jurua, Western Brazilian Amazon. *Bot J Linn Soc* 108: 213-237.
- Cayuela L, Golicher DJ, Benayas, JMR, Gonzalez-Espinosa MG, Ramirez-Marcial N. 2006. Fragmentation, disturbance and tree diversity conservation in tropical montane forests. *J Appl Ecol* 43: 1172-1181.
- Curtis JT, McIntosh RP. 1950. The interrelations of certain analytic and Synthetic phytosociological characters. *Ecology* 31: 434-455.
- Haridasan K, Rao PR. 1985-1987. *Forest Flora of Meghalaya*. Vol. 1 and 2. Bishen Singh Mahendra Pal Singh, Dehra Dun.
- Jha CS, Goparaju L, Tripathi A, Gharai B, Raghubanshi AS, Singh JS. 2005. Forest fragmentation and its impact on species diversity: an analysis using remote sensing and GIS. *Biodiv Conserv* 14: 1681-1698.
- Kadavul K, Parthasarathy N. 1999. Structure and composition of woody species in tropical semi evergreen forest of Kalrayan hills, Eastern Ghats, India. *Trop Ecol* 40 (2): 247-260.
- Kanjilal UN, Kanjilal PC, Das A, De RN, Bor NL. 1934-1940. *Flora of Assam*. 5 vols. Government Press, Shillong.
- Khan ML, Menon S, Bawa KS. 1997. Effectiveness of the protected area network in biodiversity conservation: a case study of Meghalaya, India. *For Ecol Manag* 14: 293-304.
- Khera N, Kumar A, Ram J, Tewari A. 2001. Plant biodiversity assessment in relation to disturbance in mid-elevation forest of central Himalaya, India. *Trop Ecol* 42 (1): 83-95.
- Laloo RC, Kharlukhi L, Jeeva S, Mishra BP. 2006. Status of medicinal plants in the disturbed and the undisturbed forest of Meghalaya North-east India: Population structure and regeneration efficacy of some important species. *Curr Sci* 90 (2): 225-231.
- Lalramnghinglova H. 2003. *Enthnomedicinal Plants of Mizoram*. Bishen Singh Mahendra Pal Singh, Dehradun.
- Lamb D, Erskine PD, Parrotta JA. 2005. Restoration of degraded tropical forest landscapes. *Science* 310: 1628-1932.
- Metha JP, Tewari SC, Bhandari BS. 1997. Phytosociology of woody vegetation under different management regimes in Garhwal Himalaya. *J Trop For Sci* 10: 24-34.
- Mishra BP. 2010. A study on the micro-environment, litter accumulation on forest floor and available nutrients in the soils of broad-leaved, mixed pine and pine forests at two district altitude in Meghalaya, northeast India. *Curr Sci* 90 (12): 1829-1833.
- Mishra BP, Laloo RC. 2006. A comparative analysis of vegetation and soil characteristics of montane broad-leaved, mixed pine and pine forests of northeast India. In: Trivedi PC (ed) *Advances in Plant Physiology*. I.K. International Publishing House, New Delhi.
- Mishra BP, Tripathi OP, Laloo RC. 2005. Community characteristics of a climax subtropical humid forest of Meghalaya and population structure of ten important tree species. *Trop Ecol* 46: 241-251.
- Mishra BP, Tripathi OP, Tripathi RS, Pandey HN. 2004. Effects of anthropogenic disturbance on plant diversity and community structure of a sacred grove in Meghalaya, North-east India. *Biodiv Conserv* 13: 421-436.
- Mishra BP, Tripathi RS, Tripathi OP, Pandey HN. 2003. Effects of disturbance on the regeneration of four dominant and economically important woody species in a broad-leaved subtropical humid forest of Meghalaya, Northeast India. *Curr Sci* 84 (11): 1449-1453.
- Misra R. 1968. *Ecology Workbook*. IBH Publishing Co., Calcutta.
- Mueller-Dombois D, Ellenberg H. 1974. *Aims and Methods in Vegetation Ecology*. John Wiley & Sons, New York.
- Nadakarni NM, Matelson TJ, Haber WA 1995. Structure characteristics and floristic composition of neotropical cloud forest, Montenerde, Costa Rica. *J Trop Ecol* 11: 482-495.
- Nayak DC, Srivastava R. 1995. Soil of shifting cultivated areas in Arunachal Pradesh and their suitability for land use planning. *J Indian Soc Soil Sci* 43: 246-251.
- Odum EP. 1971. *Fundamentals of Ecology*. W.B. Saunders, Philadelphia.
- Phillips EA. 1959. *Methods of Vegetation Study*. Henry Holt and Co., Inc.
- Pimm SL, Russell GJ, Gittleman JL, Brooks TM. 1995. The Future of Biodiversity. *Science* 269: 347-350.
- Pragasam A, Parthasarathy N. 2010. Landscape-level tree diversity assessment in tropical forests of southern Eastern Ghats, India. *Flora* 205: 728-737.
- Reddy MBK. 2010. Vegetation types and their influence on some secondary nutrients and micro nutrients in soils of Meghalaya forests. *Indian Forester* 1445-1459.
- Roy A, Tripathi SK, Basu SK. 2004. Formulating diversity vector for ecosystem comparison. *Ecol Mod* 179: 499-513.
- Sapkota IP, Tigabu M, Oden PC. 2010. Changes in tree species diversity and dominance across a disturbance gradient in Nepalese Sal (*Shorea robusta* Gaertn.f.) forests. *J For Res* 21: 25-32.
- Sharma CM, Ghildiyal SK, Gairola S, Suyal S. 2009. Vegetation structure, composition and diversity in relation to soil characteristics of temperate mixed broad-leaved forest along an altitudinal gradient in Garhwal Himalaya. *Indian J Sci Technol* 2 (7): 39-45.
- Singh BS, Mishra BP, Tripathi SK. 2011. Changes in phytosociological attribute and plant species diversity in secondary successional forests following stone mining in Aizawl Districts of Mizoram India. *NeBio* 2: 20-25.
- Singh BS, Mishra BP, Tripathi SK. 2012. Status of ethnomedicinally important plants sampled from different forest patches of Mizoram University campus. *NeBio* 3: 15-19.
- Singh JS, Yadav PS. 1974. Seasonal variation in composition plant biomass and net primary productivity of tropical grassland at Kurukshetra. India. *Ecol Monographs* 44: 351-376.
- Tripathi OP, Pandey HN, Tripathi RS. 2004. Distribution, community characteristics and tree population structure of subtropical pine forest of Meghalaya, Northeast India. *Intl J Ecol Environ Sci* 29: 207-214.
- Tripathi SK, Kushwaha CP, Basu SK. 2012. Application of fractal theory in assessing soil aggregates in Indian tropical ecosystems. *J For Res* 23 (3): 355-364.
- Tripathi SK, Singh RS, Upadhyay VP, Singh RP. 1989. Population structure of *Pinus roxburghii* and *Quercus leucotrichophora* forests of Almora Hills of Kumaun Himalaya. *J Tree Sci* 8: 78-80.
- Tynsong H, Tiwari BK. 2010. Plant diversity in the homegardens and their significance in the livelihoods of war khasi community of Meghalaya, Northeast India. *J Biodiv* 1 (1): 1-11.
- Tynsong H, Tiwari BK. 2011. Diversity and population characteristics of woody species in natural forests and arecanut agroforestry of south Meghalaya, Northeast India. *Trop Ecol* 52 (3): 243-252.
- Yu-Hai Y, Ya NC, We-Hong L. 2009. Relationship between soil properties and plant diversity in a desert Riparian Forest in the lower reaches of the Tarim river, Xinjiang, China. *Arid Land Res Manag* 23: 283-296.