

The bioactivity of bacterium and fungi living associate with the sponge *Reniera* sp. against multidrug-resistant *Staphylococcus aureus* and *Escherichia coli*

AGUS TRIANTO^{1,4,*}, NIRWANI¹, OKTORA SUSANTI², DIANSARI MAESAROH³, OCKY KARNA RADJASA^{1,4}

¹Department of Marine Science, Faculty of Fisheries and Marine Science, Universitas Diponegoro. Jl. Prof. Dr. Soedharto, SH., Tembalang, Semarang 50275, Central Java, Indonesia. Tel./fax. +62-24-7474698, *email: agustrianto.undip@gmail.com

² Department of Fisheries and Marine, Universitas Lampung. Jl. Prof. Dr. Sumantri Brojonegoro No.1, Bandar Lampung, Lampung, Indonesia

³Universitas Islam Negeri Sunan Ampel Surabaya. Jl. A. Yani 117 Surabaya Jl. Ahmad Yani No.117, Surabaya 60237, East Java, Indonesia

⁴Central Laboratory for Research and Services, Universitas Diponegoro. Jl. Prof. Soedarto, SH, Tembalang, Semarang 50275, Central Java, Indonesia

Manuscript received: 20 May 2019. Revision accepted: 25 July 2019.

Abstract. Trianto A, Nirwani, Susanti O, Maesaroh D, Radjasa OK. 2019. The bioactivity of bacterium and fungi living associate with the sponge *Reniera* sp. against multidrug-resistant *Staphylococcus aureus* and *Escherichia coli*. *Biodiversitas* 20: 2302-2307. The study aimed to isolate and identify the sponge-associated microorganisms producing the antibacterial substances. The sponge *Reniera* sp. was collected by hand during skin diving in Karimunjawa Islands, Indonesia. The microbial symbionts were isolated with the dilution method and screened with the overlay method against the MDR *S. aureus* and *E. coli*. The bacterium was cultured in Zobell medium, while the fungi were cultured in malt extract broth (MEB) medium. The isolates were identified based on the molecular method. A total of 46 bacteria and 43 fungi were isolated, which 7 bacteria and 20 fungi exhibited antibacterial activity against the MDR *E. coli* and *S. aureus* strains. The molecular identification revealed that the active isolates close to *Pseudoalteromonas maricaloris* (99%), *Aspergillus nomius* (96%), *Eurotium rubrum* (99%), and *Penicillium* sp. (100%). Fractionation of K.J.16.U extract gave a fraction that active to the *S. aureus* and *E. coli* strains at concentration 150 and 15 µg disk-1. The fraction K.J.16.U.1.4.4 exhibited stronger activity than that exhibited by chloramphenicol at 150 µg disk-1. The sponge *Reniera* sp. collected from Karimunjawa Islands comprise bacterial and fungal isolates produced antibacterial compounds that inhibited the growth the MDR *E. coli* and *S. aureus* strains.

Keywords: Associate microorganisms, bioactive compounds, MDR pathogen, sponges *Reniera*

INTRODUCTION

The sponge is a primitive marine organism belongs to phylum Porifera that has important ecological roles as substrate stabilizer, bio-eroding, a shelter for small organisms, and in the cycle of silicon, carbon, nitrogen, and other nutrients (Bell 2008). As a sessile and filter feeder organism, the sponge is susceptible to physical damage, predatory attacks, and pathogenic infections. Sponges have developed the specific morphology, physiology, ability to produce diverse bioactive metabolites to support their survival. Some of the secondary metabolites possessed interesting bioactivities such as antibacterial (Salim et al. 2012), antifungal (Taylor et al. 2007), antiviral, anti-inflammatory, antinociceptive (Souza et al. 2009), anticancer, and antioxidant (Trianto et al. 2011).

The *Reniera* sp. one of the sponges belongs to the Chalinidae family that well known as a rich source of the bioactive compounds, have attracted many researchers. Saito et al. (2004) isolated several bioactive compounds belong to reniera mycin, mimosamycin, and renierone that have bioactivity as antiproliferative against several cancer cells. Lunder et al. (2012) published the polymeric 3-alkylpyridinium salts, isolated from the sponge *Reniera sarai*, a unique water-soluble compound that has a cardiotoxic effect. Urda et al. (2018) exhibited the

bioactive compounds from *Reniera* sponge called Njoamines, a member of the alkaloid family contains a tricyclic nitrogenated core.

The surface or the inner parts of the sponges rich in nutrients than those present in seawater and sediments that suitable habitats for microorganisms. Some of these microorganisms are used as food while others live in symbiosis with the sponge as their host. Although the interaction between sponges and microorganisms is not yet clearly understood, there is an exchange of nutrients and secondary metabolites, which suggests that the survival of the sponge presumably depends on the positive host-symbiont relationship. This kind of interaction may contribute to the nutrient acquisition and production of secondary metabolites in responsibility to protect the sponge body from microbial infection (Engel et al. 2002). Some of the bacteria living symbiotically with the sponge produce bioactive compounds (Kikuchi et al. 2018). Marine sponges are also rich in fungal symbionts that produce antibacterial and fungal compounds (Trianto et al. 2017a, b). Marine microbes, both living freely and in association with the sponges, have been known to produce various bioactive compounds. The purpose of this work is to assess antibacterial the potential of bacteria and fungi isolated from sponge *Reniera* sp. collected from

Karimunjawa Islands, Indonesia against two multi-resistant resistance bacteria (MDR).

MATERIALS AND METHODS

Sponge collection and isolation of sponge microbial symbionts

The sponge *Reniera* sp. was collected from Karimunjawa Islands, Jepara, Central Java, Indonesia, from the depth of 2-5 m by skin diving method. Each sponge specimen was put into a sterile zip-lock plastic bag and stored in a cool box until the isolation process. The specimens were washed with the sterile seawater prior to microbial isolation. The fungal isolates were isolated from the sponge by tapping method. A small part (ca. 1 cm x 1 cm x 0.3 mm) of was cut from each specimen, and the sponge part was put onto the malt extract agar (MEA) media surface. While the microbial symbionts were isolated using the serial dilution method. The tissue solutions were spread on Zobell marine agar plates (Radjasa et al. 2007). Identification of the sponge was based on the in situ picture and spicule analysis (Hooper 2003; Kang et al. 2013).

Screening for antibacterial activity

Both of the bacterial and the fungal isolates were screened for the antibacterial activity against the pathogenic bacteria MRSA and MDR *E. coli* using the modified overlay method described by Anand et al. (2006). The bacterial symbionts were spotted on Zobell marine agar medium and incubated for three days, while the fungal strains were inoculated in the MEA medium and incubated for four days. The fresh culture of the pathogens was mixed with soft agar medium (1% v/v) and overlaid over the agar surface of the previous microbial symbiont cultures. The plates were then incubated at 37°C for 24 h. The antibacterial activity was defined by the presence of clear zones around the microbial symbiont.

Gram test

The bacterial Gram identification was carried out using the KOH 3% string and Gram staining tests. KOH 3% string test was done according to Ali et al. (2015); Dash and Payyappilli (2016), a loopful of the bacterial colony from the culture plate was emulsified over glass slide in the suspension of 3% KOH. Gram staining test was done according to Ayuningrum et al. (2017), using Gram's staining solution A (crystal violet), solution B (Lugol iodine), solution C (alcohol) and solution D (safranin).

Mass culture and extraction of microbial symbionts

The isolates with the strongest activity were chosen for further studies. The most active isolates, a bacterium, and three fungal isolates were cultured for the production of the bioactive compounds. The bacterial inoculum was prepared in 5-mL Zobell broth medium on a rotary shaker at room temperature, after 24 h, the inoculum was serially

transferred into larger volumes (10, 100, and 1000 mL). The fungi were cultured in malt extract broth (MEB) medium at room temperature for seven days (Radjasa & Sabdono 2003). The microbial mass was separated from the culture medium for extraction of the bioactive substances. The bacterial culture was centrifuged at 1000 rpm for 10 min, the supernatant was discarded, and the pellet was soaked in MeOH for 24 h. The fungal mycelia were separated from the culture medium by filtration with a paper filter; then the mycelia were soaked in MeOH for 24 h. The solution was filtered and concentrated with the rotary evaporator in vacuo to obtain the crude extracts. The extracts were partitioned using a solvent mixture of ethyl acetate and H₂O solvents to provide the organic and water fractions.

Antibacterial test of the extract

The antibacterial test was conducted by the disk diffusion method. The extracts were dissolved in ethyl acetate to prepare the stock solutions with the concentration 1.5 µg µL⁻¹, then 10 µL of the stock solutions was impregnated into a sterile paper disk with diameter 5 mm in to give the final concentration at 15 µg disk⁻¹. The paper disks were placed on the nutrient agar medium surface that was previously inoculated with the pathogenic bacteria. The standard antibiotic (chloramphenicol, 15 µg disk⁻¹) was used in this test. The antibacterial activity was defined based on the inhibition zones around the paper disk (Ozdemir et al. 2006).

Molecular identification

The DNA extraction was carried out using the Chelex method (Walsh et al. 1991). Selected colonies were inoculated in 50-100 µL ddH₂O and 1 mL of 0.5% saponin in phosphate buffer solution (PBS) 1× (stored overnight). The mixture was centrifuged at 12000 rpm for 10 min (1 rpm = 1/60 Hz), and then the supernatant was discarded. Then, 100 µL ddH₂O and 50 µL of 20% Chelex 100 were added to the final solution, and the solution was boiled for 10 min and vortexed once for 5 min. The mixture was centrifuged at 12000 rpm for 10 min and stored at -20°C.

The bacterial DNA that was used for 16S rRNA gene sequencing was amplified by polymerase chain reaction (PCR) using the universal primers 27F (5'AGAGTTTGATCMTGGCTCAG-3') and 1492R (5'TACGGTTAACCTTG TTACGACTT-3'). The fungal DNA that was used for 5.8S rDNA gene sequencing was amplified by PCR with the internal transcribed spacer (ITS) region ITS 1 (5'TCCGTAGGTGAA CCTGCGG-3') and the primer ITS4 (5'TCCTCCGCTTATTGATATGC-3').

The DNA sequencing was conducted at Macrogen Laboratory, Korea. The results of DNA sequences were preliminarily aligned with ClustalW Multiple Sequence Alignment. Phylogenetic analysis was performed using MEGA 6. The phylogenetic trees were determined by the neighbor-joining method using Kimura's two-parameter model.

RESULTS AND DISCUSSION

Isolation the microbial and screening for antibacterial activity of the isolates

A total of 46 bacteria and 43 fungi were isolated from the sponge *Reniera* sp. Among the isolates, 5 bacteria inhibited *S. aureus* and *E. coli*, and 3 of the active isolates inhibited both pathogenic bacteria. Whilst, 19 and 7 fungal isolates inhibited *S. aureus* and *E. coli*, respectively. Among the active fungal isolates, 6 of them inhibited both bacteria (Figure 1). The preliminary test revealed that 23.6% of the strains were capable of inhibiting the MDR pathogens MRSA and *E. coli*. The antibacterial test of the isolates extract showed one bacterium along with three fungal isolates having strong activity against the antagonist bacteria.

The *Reniera* sponge is known to have symbiotic microbes either bacteria or fungi. Sponge *Reniera* sp. collected from Panjang Island, Jepara Central Java contains five associated bacteria that one of the bacteria is *Chromohalobacter salexigens* strain DSM 3043 active against *P. aeruginosa*, *E. cloacae*, *A. baumannii*, and *S. aureus* (Asagabaldan 2017). The microbial community plays important roles in sponge survival and adaptation to environmental change. The microorganisms serve the food supply of the sponge both internal exchange and with the environment (Kiran et al. 2018). Bacteria also produced bioactive compounds that protect the sponge from biofouling organisms (Aguila-Ramírez et al. 2014; Satheesh et al. 2016).

Bioactivity of bacterial biomass and fungal mycelia extract

The most active isolates were cultured in higher mass to provide the extract for further study. The extraction of the bacterial biomass and fungal mycelia yielded methanolic extract with concentrations of 5.3-18.7%. The methanol extraction yielded crude extract that was partitioned with H₂O and ethyl acetate before conducting the antibacterial assay. The antibacterial screening results are shown in Table 1.




Figure 1. The number of total and active bacterial and fungal isolates of the sponge *Reniera* sp.

Table 1. Bioactivity of ethyl acetate extract of the bacterial biomass and fungal mycelia obtained from 1liter batch culture

Isolat	Bacterial/ Mycelial weight (g)	Extract weight (mg)	Extract concentration (%)	The diameter of the inhibition zone (mm)	
				<i>E. coli</i>	MRSA
SKSBT.0.3.29	6.6	348.4	5.3	2.3	7.2
SKSJT. J. 16 H	17.9	3340.0	18.7	10.3	10.6
SKSJT. J. 16 U	31.0	3280.0	10.6	10.7	9.6
SKSJT. J.19	9.4	1300.0	13.8	7.4	2.5

Note: The diameter of the inhibition zone of (+) control chloramphenicol: 14.83 and 13.95 mm against *E. coli* and *S. aureus*, respectively. (-) control ethyl acetate and H₂O fraction did not show activity (0.00 mm).

Table 3. Identification of fungal strains isolated from *Reniera* sp. Samples were based on BLAST analysis of the internal transcribed spacer (ITS) region

Strain	Sequence length (nt)	Related cultivated strain (BLAST)	Acc. no	Similarity (%)
SKSBT.0.3.29	1165	<i>Pseudoalteromonas maricaloris</i> strain KMM636	NR 025009.1	99%
K.J.16.U	1062	<i>Aspergillus nomius</i> strain SGE19	JN709035.1	96%
K.J.16.H	545	<i>Eurotium rubrum</i> strain SGE27	JX232274.1	99%
K.J.19	983	<i>Penicillium</i> sp. DY115-21-10-M23	KF411603.1	100%




Figure 2. Phylogenetic tree of bacterial isolate




Figure 3. Phylogenetic tree of fungal isolates

Marine bacteria and fungi are well known as potential sources of natural products. For example, *Pseudoalteromonas* sp. reported producing compound pentabromopseudilin, a promising anti-MRSA and myosin ATPase inhibition with activities at a level comparable to those of commercial drugs (Kum et al. 2017). Other researches showed that *Pseudoalteromonas* sp. living

associated with sponge *Halisarca ectofibrosa* produce compounds that inhibit the growth of *Bacillus subtilis* and *Vibrio anguillarum* (Rungprom et al. 2008; Schinke et al. 2017). A review paper by Xu et al (2015) describes broadly the potential of marine fungi as sources bioactive compounds. The review showed there are 105 marine fungal strains used for the isolation of antibacterial or

antifungal compounds. Mangrove and sponges are the most rich sources of fungi, while *Aspergillus* and *Penicillium* are the most common fungi in the marine ecosystem. Other review depicted the high potential of marine fungi especially the genus *Aspergillus* of antibacterial and anticancer (Bladt et al. 2013).

Identification of the isolates

The ethyl acetate fraction of K.J.16.U was selected. Molecular identification of the active bacterial and fungal symbionts based on 16S rRNA and 5.8S rDNA revealed that the active strains SKSBT.0.3.29, K.J.16.U, K.J.16.H, and K.J.19 were closely related to *Pseudoalteromonas maricaloris*, *Aspergillus nomius*, *Eurotium rubrum*, and *Penicillium* sp., respectively for bioassay-guided fractionation because of its high antibacterial activity, which yielded six fractions (K.J.16.U.1.4.1-6) (Table 2). The results obtained from BLAST homology of the bacterial and fungal symbionts of *Reniera* sp. are described by phylogenetic trees (Figures 2 and 3).

The molecular-based identification indicated that the active isolate is *Pseudoalteromonas maricaloris* with the BLAST homology 99% similarity. While the fungal isolates identified as *Aspergillus nomius* strain SGE19, *Eurotium rubrum* strain SGE27, and *Penicillium* sp. DY115-21-10-M23 with BLAST homology 99% similarity 96%, 99%, and 100% similarity, respectively (Table 2). Genus *Pseudoalteromonas* are heterotrophic bacteria include are essential components of the marine environment and have diverse habitats including coastal and open water areas, bottom sediments, and living associated with marine plants and animals. *P. maricaloris* has been isolated from a sponge, *Fascaplysinopsis reticulata*, collected from Australia. It has also been reported having antibacterial activity against *S. aureus*, *E. coli*, *Proteus vulgaris*, *Enterococcus faecium*, and *Bacillus subtilis* and also cytotoxic activity against tumor cells (Ivanova et al. 2002; Radjasa et al. 2007). *Pseudoalteromonas* also produces extracellular antibacterial compounds that inhibit the *Vibrio parahaemolyticus* causing acute hepatopancreatic necrosis disease (Wang et al. 2018). Asagabaldan et al. (2017) isolated bacterial strains from the sponge belonging to the genus *Haliclona* (Reneira) that were active against *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Acinetobacter baumannii*, and *Enterobacter cloacae*.

Penicillium sp., *Aspergillus* sp., and *Eurotium* sp. have been isolated from the sponge *Tethya aurantium* collected from the Istrian Peninsula, Croatia, which possesses the secondary metabolite mycotoxin that exhibited antineoplastic properties. *A. nomius* is reported as an insect parasite and a producer aflatoxin B1, B2, G1, and G2, mycotoxins, that toxic to humans, animals, and insects (Calderari et al. 2013; Afonso et al. 2017). However, *A. nomius* has also known as lipase producer that potentially used for the production of fatty acid methyl ester in the biodiesel industry (Rakchai et al. 2016). *Eurotium rubrum* commonly found in grain products, poultry feed, bakery products, dried fruits, spices, soil, hypersaline waters, and Dead Sea soil (Butinar et al. 2005). These *Eurotium*

reported to producing diketopiperazine compounds that potential in the cosmetic industry by inhibited melanogenesis (Kamauchi et al. 2016). The *E. rubrum* has also reported producing a metabolite called flavoglaucin, auroglaucin, isotetrahydroauroglaucin, eochinulins A and B, echinulin, preechinulin, nechinulin E, epihevadride, and questin. The compounds have activities as anticancer, antioxidant, antibacterial, cytoprotection against peroxynitrite, hepatotoxic (Slack et al. 2009). *Penicillium* sp. is well-known for its ability to produce secondary metabolites with diverse biological effects. *Penicillium* sp. has been reported contains Peniopyranone, a bioactive compound inhibits acetylcholinesterase, that potential for neurodegenerative diseases such as Alzheimer's dementia and Parkinson's disease (Li et al. 2017). *Penicillium* sp. also well-known as a source of prenylated indole alkaloids having interesting biological activities such as antitumor, anthelmintic, calmodulin inhibitory, and antibacterial (Yang et al. 2018).

To conclude, the sponge *Reniera* sp. collected from Karimunjawa Islands comprise bacterial and fungal isolates produced antibacterial compounds that inhibited the growth the MDR *E. coli* and *S. aureus* strains. The fungus *A. nomius* has potential as a source of antibacterial compounds that having stronger activity against MRSA and *E. coli*.

ACKNOWLEDGMENTS

This research was partially supported by The Ministry of Research, Technology, and Higher Education through Post Graduate Scheme. We thank USAID for supporting this research through PEER grant.

REFERENCES

- Afonso E, Raquel C, Valadares L, Peoporine N, Santos F, Tallarico M. 2017. Aflatoxins produced by *Aspergillus nomius* ASR3, a pathogen isolated from the leaf-cutter ant *Attasexdens rubropilosa*. Rev Brasileira Farmacog 27: 529-532.
- Aguila-Ramírez RN, Hernández-Guerrero CJ, González-Acosta B, Id-Daoud G, Hewitt S, Pope J. 2014. Antifouling activity of symbiotic bacteria from sponge *Aplysina gerardogreeni*. Intl Biodeterior Biodegrad 90: 64-70.
- Ali F, Bahadar S, Rahman KU, Khan I, Zeb A, Khan J, Daud M, Hassan A, Rabbani NK, Khattak AA, Jadoon FK, Hayat. 2015. Isolation and identification of lipase producing bacteria from coal mines of Darra Adam Khel, Kyber Pakhtoonkha, Pakistan. Int J Biosci 7 (2): 187-191.
- Anand TP, Bhat AW, Shouche YS, Roy U, Siddharth J, Sarma SP. 2006. Antimicrobial activity of marine bacteria associated with sponges from the waters off the coast of southeast India. Microbiol Res 16: 252-262.
- Asagabaldan MA, Ayuningrum D, Kristiana R, Sabdono A, Radjasa OK, Trianto A. 2017. Identification and antibacterial activity of bacteria isolated from marine sponge *Haliclona (Reniera)* sp. against multi-drug resistant human pathogen. IOP Conf Ser: Earth Environ Sci. 55: 012019. DOI: 10.1088/1755-1315/55/1/012019
- Ayuningrum D, Kristiana R, Asagabaldan MA, Sabdono A, Radjasa OK, Nuryadi H, Trianto A. 2016. Isolation, characterisation and antagonistic activity of bacteria symbionts hard coral *Pavona* sp. isolated from Panjang Island, Jepara against infectious Multi-drug Resistant (MDR) bacteria. IOP Conf Ser: Earth Environ Sci. 55: 012019. DOI: 10.1088/1755-1315/55/1/012019

- Bell JJ. 2008. The functional roles of marine sponges. *Estuar Coast Shelf Sci* 79 (3): 341-353.
- Bladt TT, Frisvad JC, Knudsen PB, Thomas Ostenfeld Larsen. 2013. Anticancer and antifungal compounds from *Aspergillus*, *Penicillium*, and other filamentous fungi. *Molecules* 18: 11338-11376.
- Butinar L, Zalar P, Frisvad JC, Gunde-Cimerman N. 2005. The genus *Eurotium* members of indigenous fungal community in hypersaline waters of salterns. *FEMS Microbiol Ecol* 51: 155-166
- Calderari TO, Iamanaka BT, Frisvad JC, Pitt JI, Sartori D, Luiz J. 2013. The biodiversity of *Aspergillus* section Flavi in brazil nuts: From rainforest to consumer. *Intl J Food Microbiol* 160: 267-272.
- Dash C, Payyappilli RJ. 2016. KOH string and Vancomycin susceptibility test as an alternative method to Gram staining. *J Intl Med Dent* 3 (2): 88-90.
- Engel S, Jensen PR, Fenical W. 2002. Chemical ecology of marine microbial defense. *J Chem Ecol* 28: 1971-1985.
- Hooper JNA. 2003. Sponguide: Guide to sponge collection and identification, version 2003. Queensland Museum, Australia.
- Ivanova EP, Shevchenko LS, Sawabe T, Lysenko AM, Svetashev VI, Gorshkova NM, Satomi M, Christen R, Mikhailov VV. 2002. *Pseudoalteromonas maricaloris* sp. nov. isolated from an Australian sponge and reclassification of *Pseudoalteromonas aurantia* NCIMB 2033 as *Pseudoalteromonas flavipulchra* sp. nov. *Intl J Syst Evol Microbiol* 52: 32-71.
- Kamauchi H, Kinoshita K, Sugita T, Koyama K. 2016. Conditional changes enhanced production of bioactive metabolites of marine-derived fungus *Eurotium rubrum*. *Bioorg Med Chem Lett* 26: 4911-4914.
- Kang DW, Kyung JL, Chung JS. 2013. Two new marine sponges of the genus *Haliclona* (Haplosclerida: Chalinidae) from Korea. *Anim Syst Evol Divers* 29: 51-55.
- Kikuchi S, Okada K, Cho Y, Yoshida S, Kwon E, Yotsu-Yamashita M. 2018. Isolation and structure determination of lysiformine from bacteria associated with marine sponge *Halichondria okadae*. *Tetrahedron* 74 (27): 3742-3747.
- Kiran GS, Sekar S, Ramasamy P, Thinesh T, Hassan S, Lipton AN. 2017. Marine sponge microbial association: Towards disclosing unique symbiotic interactions. *Mar Environ Res* 140: 169-179
- Kum D-Y, Nazari M, McPhail KL, Cooper CS, Suyama TL. 2017. Two-step total synthesis of an anti-MRSA and myosin-inhibiting marine natural product pentabromopseudinil via Suzuki-Miyaura coupling of a MIDA boronate ester. *Tetrahedron Lett* 58: 3374-3376.
- Li XS, Tan MH, Lu MM, Liu CX, Guo ZY, Zhang L. 2017. New γ -pyranone and nucleoside derivatives from *Penicillium* sp. *Phytoch Lett* 20: 285-288.
- Lunder M, Drevnšek G, Hawlina S, Sepčić K, Žiberna L. 2012. Cardiovascular effects induced by polymeric 3-alkylpyridinium salts from the marine sponge *Reniera sarai*. *Toxicon* 60 (6): 1041-1048.
- Ozdemir G, Karabay NU, Dalay MC, Pazarbasi B. 2006. Antimicrobial activities of the volatile component and various extracts of *Dictyopteria membranacea* and *Cystoseira barbata* from the coast of Izmir, Turkey. *Pharm Biol* 44: 183-188.
- Radjasa OK, Sabdono A. 2003. Screening of secondary metabolite-producing bacteria associated with corals using 16S rDNA-based approach. *J Coast Develop* 7: 11-19.
- Radjasa OK, Salasia SIO, Sabdono A, Weise J, Imhoff JF, Lammler C, Risk MJ. 2007. Antibacterial activity of marine bacterium *Pseudomonas* sp. associated with soft coral *Sinularia polydactyla* against *Streptococcus equi* subsp. zoepidemicus. *Intl J Pharmacol* 3: 170-174.
- Rakchai N, Kittikun A, Zimmermann W. 2016. The production of immobilized whole-cell lipase from *Aspergillus nomius* ST57 and the enhancement of the synthesis of fatty acid methyl esters using a two-step reaction. *J Mol Catal B Enzym* 133: 10-13.
- Rungprom W, Siwu ERO, Lambert LK, Dechsakulwatana C, Barden MC, Kokpol U, Blanchfield JT, Kita M, Garson MJ. 2008. Cyclic tetrapeptides from marine bacteria associated with the seaweed *Digenea* sp. and the sponge *Halisarca ectofibrosa*. *Tetrahedron* 64: 3147-3152
- Saito N, Tanaka C, Koizumi YI, Suwanborirux K, Amnuoyopol S, Pummangura S. 2004. Chemistry of *Reniera* mycins. Part 6: Transformation of *Reniera* mycin M into jorumycin and *Reniera* mycin J including oxidative degradation products, mimosamycin, renierone, and renierol acetate. *Tetrahedron* 60 (17): 3873-3881.
- Salim AA, Khalil ZG, Capon RJ. 2012. Structural and stereochemical investigations into bromotyrosine-derived metabolites from southern Australian marine sponges *Pseudoceratina* spp. *Tetrahedron* 6: 9802-9807.
- Satheesh S, Ba-akdah MA, Al-Sofyani AA. 2016. Natural antifouling compound production by microbes associated with marine macroorganisms — A review. *Electron J Biotechnol* 21: 26-35.
- Schinke C, Martins T, Queiroz SCN, Melo IS, Reyes FGR. 2017. Antibacterial compounds from marine bacteria, 2010-2015. *J Nat Prod* 80: 1215-1228.
- Slack GJ, Puniani E, Frisvad JC, Samson RA, Miller JD, Peterson SW. 2009. Secondary metabolites from *Eurotium* species, *Aspergillus calidoustus*, and *A. insuetus* common in Canadian homes with a review of their chemistry and biological activities. 113: 480-490.
- Souza ET, de Lira DP, de Queiroz AC, da Silva DJC, de Aquino AB. 2009. The antinociceptive and anti-inflammatory activities of caulerpin, a bisindole alkaloid isolated from seaweeds of the genus *Caulerpa*. *Mar Drugs* 7: 689-704.
- Taylor MW, Radax R, Steger D, Wagner M. 2007. Sponge-associated microorganisms: Evolution, ecology, and biotechnological potential. *Microbiol Mol Biol Rev* 71: 295-347.
- Trianto A, Hermawan I, de Voogd NJ, Tanaka J. 2011. Halioxepine, a new meroditerpene from an Indonesian sponge *Haliclona* sp. *Chem Pharm Bul* 59: 1311-1313.
- Trianto A, Radjasa OK, Pribadi R, Widyaningsih S, Witriansyah K, Yusidharta I, Wiratno, Riniatsih I. 2017a. Isolation and identification of sponge-associated fungus producing anti-multidrug-resistant (MDR) bacterial substances. *Res J Pharm Biol Chem Sci* 8: 63.
- Trianto A, Sabdono A, Rochaddi B, Triningsih D W. 2017b. Exploration of marine sponges-associated fungi producing antifungal compounds. *Asian J Microbiol Biotech Environ Sci* 19: 588-593.
- Urda C, Pérez M, Rodríguez J, Fernández R, Jiménez C, Cuevas C. 2018. Njaoamine I, a cytotoxic polycyclic alkaloid from the Haplosclerida sponge *Haliclona (Reniera)* sp. *Tetrahedron Lett* 59 (26): 2577-1580.
- Walsh PS, Metzger DA, Higuchi R. 1991. Biotechniques 30th Anniversary gem: Chelex 100 as a medium for simple extraction of DNA for PCR-based typing from forensic material. *Biotechniques* 10: 506-513.
- Wang H, Wang C, Tang Y, Sun B, Huang J, Song X. 2018. *Pseudoalteromonas* probiotics as potential biocontrol agents improve the survival of *Penaeus vannamei* challenged with acute hepatopancreatic necrosis disease (AHPND)-causing *Vibrio parahaemolyticus*. *Aquaculture* 494: 30-36.
- Xu L, Meng W, Cao C, Wang J, Shan W, Wang Q. 2015. Antibacterial and antifungal compounds from marine fungi. *Mar Drugs* 13: 3479-3513.
- Yang B, Tao H, Lin X, Wang J, Liao S, Dong J. 2018. Prenylated indole alkaloids and chromone derivatives from the fungus *Penicillium* sp. SCSIO041218. *Tetrahedron* 74 (1): 77-82.