

## Short Communication:

# Single nucleotide polymorphism in C-type lysozyme gene and its correlation with *Aeromonas hydrophila* resistance in African catfish *Clarias gariepinus*

HASAN NASRULLAH<sup>1</sup>, YANTI INNEKE NABABAN<sup>1</sup>, IKA SAFITRI<sup>1</sup>, DWI HANI YANTI<sup>2</sup>, SRI NURYATI<sup>1</sup>, MUHAMMAD ZAIRIN JUNIOR<sup>1</sup>, ALIMUDDIN ALIMUDDIN<sup>1,\*</sup>

<sup>1</sup>Department of Aquaculture, Faculty of Fisheries and Marine Sciences, Institut Pertanian Bogor. Jl. Agatis, IPB Dramaga Campus, Bogor 16680, West Java, Indonesia. Tel. +62-251-8622755, Fax.: +62-251-8622941, \*email: alimuddin@apps.ipb.ac.id

<sup>2</sup>National Center for Freshwater Aquaculture. Jl. Selabintana No. 37, Sukabumi 43114, West Java, Indonesia

Manuscript received: 23 October 2019. Revision accepted: 28 December 2019.

**Abstract.** Nasrullah N, Nababan YI, Safitri I, Yanti DH, Nuryati S, Zairin Junior M, Alimuddin A. 2020. Short Communication: Single nucleotide polymorphism in C-type lysozyme gene and its correlation with *Aeromonas hydrophila* resistance in African catfish *Clarias gariepinus*. *Biodiversitas* 21: 311-317. The chicken-type lysozyme (LYSC) gene has been demonstrated to play important roles in the fish protection system against bacterial infection. In this present study, we aimed to identify the single nucleotide polymorphism (SNP) within the LYSC gene of African catfish *Clarias gariepinus* and its possible association with *Aeromonas hydrophila* resistance, a major pathogenic bacterium in African catfish. The gDNA of the African catfish LYSC coding sequence was 1559 bp long, comprising of four exons and three introns. Six SNPs were identified in African catfish LYSC, namely SNP1-6. After the *A. hydrophila* challenge, we regarded the surviving individuals after the infection as the resistant group and the dead fish as the susceptible group. High-resolution melting (HRM) analysis on SNP2 revealed that the allele frequencies of TT, CC, and TC were of 27.78%, 5.55%, and 66.67% in the resistant group, while the frequencies of TT, CC, and TC were 16.67%, 27.78%, and 55.55% in the susceptible group, respectively. The expression of LYSC and other immune-genes in the resistant group was also higher in the liver, head kidney, and spleen. These results indicated that the LYSC gene might play an essential role in bacterial resistance, and the SNP2 within the LYSC gene may be associated with the resistance to *A. hydrophila* in African catfish.

**Keywords:** *Aeromonas hydrophila*, African catfish, disease resistance, lysozyme gene, SNP

## INTRODUCTION

The African catfish (*Clarias gariepinus* Bloch, 1794) is an important aquaculture species in the world that mainly produced in African and Asian countries, including Indonesia (Ekasari et al. 2016; Dauda et al. 2018). The African catfish production has increased significantly through intensive fish culture practices, thus consequently exposes the cultured fish to common stressors and diseases (Shoko et al. 2016; Yarahmadi et al. 2016). One of the most commonly occurred diseases in African catfish is motile aeromonad septicemia (MAS) caused by *Aeromonas hydrophila* infection. This Gram-negative bacterium has caused high mortality in African catfish and other freshwater species, leading to substantial economic loss (Angka et al. 1995; Hanson et al. 2014). Most cases of MAS can be correlated to various factors such as infection, and environmental stress, and sometimes associated with nutritional stress and handling stress (Hanson et al. 2014). Recently, our research has begun to focus on breeding African catfish resistant to *A. hydrophila*. However, little information is available about genes related to *A. hydrophila* resistance in African catfish.

Lysozyme has crucial roles in fish early defense system

and innate immunity. This hydrolase enzyme catalyzes the hydrolysis of  $\beta$ -(1,4)-glycosidic bonds between the N-acetyl glucosamine and N-acetyl muramic acid in the peptidoglycan layer of bacterial cell walls, thus possess bacteriolytic activity against Gram-positive and Gram-negative (Saurabh dan Sahoo 2008). Generally, among the vertebrate and teleost fish, lysozyme is mostly found in two types: chicken-type (LYSC) and goose-type (LYSG) (Callewaert dan Michiels 2010; Buonocore et al. 2014). Both lysozymes gene has been demonstrated to play essential roles as the immune defense in many fish species (Mohanty dan Sahoo 2010; Pridgeon et al. 2013; Wei et al. 2014; Di Falco et al. 2017). The involvement of the LYSC in the protection against bacterial infection has been reported in fish species, such as resistant families of Atlantic salmon against *Piscirickettsia salmonis* infection (Dettleff et al. 2015), channel catfish against *Edwardsiella ictaluri* (Wang et al. 2013), Darby's sturgeon and Qihe crucian carp against *A. hydrophila* (Wang et al. 2016; Zhang et al. 2018). Moreover, the single nucleotide polymorphisms (SNPs) in LYSC gene have been characterized in several fish species, such as Asian seabass (Fu et al. 2013) and Japanese flounder (Liu et al. 2017) and identified as potential markers for bacterial-resistance

in fish. However, such studies in African catfish are still limited. In this present study, we aimed to identify the single nucleotide polymorphism (SNP) within the LYSC gene of African catfish and studied its possible association with *Aeromonas hydrophila* resistance.

## MATERIALS AND METHOD

### Isolation of LYSC Genomic DNA sequences

To obtain the genomic sequence of the LYSC gene, primers were designed to amplify the full length of the coding sequence of LYSC mRNA (Genbank accession number: MK344777.1). The forward primer was located from the start codon, and the reverse primer sequence was ended in the stop codon of the coding sequence. *Bam*HI site was added in the forward primer and *Nde*I site in the reverse primer for future cloning purposes (Table 1). The genomic DNA was extracted from the fin using Genra® Purgene® Tissue kit (Qiagen, USA) according to the kit manual. PCR amplification was performed in a 50 µL of reaction volume consisting of 50 ng of genomic DNA, 10 µM of each primer, and 2× MyTaq™ HS DNA polymerase mix (Bioline, UK). The amplification program consisted of the following steps: 95 °C for 2 min followed by 35 cycles of 95 °C for 30 s, 60 °C for 30 s, and 72 °C for 1 min 30 s, then a final step of 72 °C for 5 min. The PCR products were purified using Wizard® Genomic DNA Purification kit (Promega, USA) and then sequenced using 3500 Genetic Analyzer (Applied Biosystem, USA). The African catfish LYSC gDNA sequence then compared to other fish species LYSC sequences available at the Genbank database.

### Fish and bacterial infection

African catfish (6.00±0.56 g, n = 1,000) were obtained from the National Centre for Freshwater Aquaculture, Sukabumi, West-Java, Indonesia. The fish were acclimated for one week in a 1,050 L fiber tank. Fish were fed twice with commercial pellets. Feed and fecal wastes were removed daily. No clinical signs of bacterial disease were observed during acclimatization. Pathogenic *A. hydrophila*

was obtained from our lab culture collection. The isolate was characterized using biochemical tests, and the lethal dose 70 (LD70) of the bacteria was determined before the experiment. *A. hydrophila* was diluted in phosphate buffer saline (PBS) solution in a final concentration of 10<sup>8</sup> CFU mL<sup>-1</sup> based on the LD70 concentration. Before the *A. hydrophila* challenge, the fish was anesthetized with 75 mg L<sup>-1</sup> MS222. Two hundred fish were injected with 0.1 mL of pathogenic *A. hydrophila* at LD70 concentration intramuscularly. Sixty fish were injected by 0.1 mL PBS as a control group. The fish were reared for 24 h in the 60 L glass tanks with 20 fish/tank density. After 24 h, we regarded the surviving individuals after the infection as a resistant group and the dead fish as the susceptible group.

### qPCR and SNP analysis

To identify the reference genotype in the LYSC gene, we amplify the gDNA from seven resistant fish, and five susceptible fish then sequenced the PCR products. The LYSC and other immune-related genes: goose-type lysozyme (LYSG), hepcidin (HAMP), and interleukin-1β (IL1B) mRNA expression of the resistant fish were analyzed at 0 h and 24 h after bacterial infection to investigate the immunological roles of the gene. Total RNA was extracted from the liver, head-kidney, and spleen of the resistant fish (n= 4) and analyzed using the qPCR method. The RNA extraction, cDNA synthesis, and qPCR analysis were conducted according to our previous research (Nasrullah et al. 2019). The β-actin gene was used for normalization and PBS control at 0 h as the expression calibrator. The expression data were analyzed using the 2<sup>-ΔΔCT</sup> method (Livak dan Schmittgen 2001).

**Table 1.** Mortality of *C. gariepinus* 24 h after challenge with *A. hydrophila*

Treatment	Number of challenged individuals	Survived	Dead	% Mortality
<i>A. hydrophila</i> infection	200	49	151	75.5
PBS control	60	60	0	0

**Table 2.** The primer used in this study

Primer name	Sequence (5' - 3')	Ta (°C)	Application	Reference
LYSC F	AAAGGATCCATGAAGGCTTTGGTGTCTT	60	gDNA amplification of LYC coding sequence	MK344777.1
LYSC R	CGATCATATGTTATTAACCTCCACAGCC			
SNPLYSC F	TTCCATAGTGCTAACGGGTGC	60	SNP Genotyping by HRM analysis	<i>This study</i>
SNPLYSC R	GCACACTCAACAGCTTTCGC			
qLYSC F	CGGTATGATCGGTGTGAGCTGG	60	qPCR analysis of LYSC	Nasrullah et al. (2019)
qLYSC R	CGGTCTGGGCGTTGGTATTGA			
qLYSG F	CCTAACTGGCCCAAAGAGCA	60	qPCR analysis of LYSG	Nasrullah et al. (2019)
qLYSG R	CCATACCCTCGTATGTGCGG			
qHAMP F	TATCGTCATCATCGCGTGCA	60	qPCR analysis of HAMP	MK344776.1
qHAMP R	TCCTTCGCCAATGATGCAGT			
qIL1B F	TGCACTGAATCCAAGAGCTACAGC	60	qPCR analysis of IL1B	MH341527.1
qIL1B R	CCACCTTTCAGAGTGAATGCCAGC			
qACTB F	ACCGGAGTCCATCACAATACCAGT	60	Internal control, expression normalization	Nasrullah et al. (2019)
qACTB R	GAGCTGCGTGTGGCCCTGAG			

Note: Restriction sites for future cloning purposes are written in italic

To examine the association of the SNP with the *A. hydrophila* resistance, we collected fin clips from 36 individuals from each resistant and susceptible group. The potential SNPs were genotyped using qPCR-high resolution melting (HRM) analysis (Reed et al. 2007). The primer for the analysis was designed based on the SNP2 as the potential marker. The HRM analysis was performed in 20  $\mu$ L of the final volume consisting of 4  $\mu$ L gDNA (1 ng), 0.8  $\mu$ L of each primer (10 mM), 10  $\mu$ L SensiFAST™ HRM kit (Bioline, UK), and 4.4  $\mu$ L nucleases free water. The HRM analysis conducted in Rotorgene6000 (Corrbet, USA) machine using the following program: 95 °C for 5 min followed by 40 cycles of 95 °C for 15 s, 60 °C for 10 s, and 72 °C for 10 s, then followed by an increasing temperature 72-92°C. Melt curves were generated in the HRM analysis module of RotorGeneQ software (Corrbet, USA). The normalized regions were set from 78-79 °C and 80.93-81.99 °C. The samples were assigned to a given genotype by examining the normalized melt curve plots in comparison to the reference genotypes at a 90% confidence threshold.

## RESULTS AND DISCUSSION

### Genomic structure and mRNA expression

The genomic sequence of African catfish LYSC gene from the transcriptional start site to the transcriptional end site was 1559 bp long, comprising of four exons (127, 159, 76 and 67 bp, respectively) and three introns (595,104, and 431 bp, respectively), as shown in Figures 1 and 2. This result is consistent with other fish species (Hikima et al. 2000; Fu et al. 2013). A phylogenetic tree (Figure S1) was constructed using the African catfish LYSC gDNA and other fish species using the neighbor-joining method. The African catfish LYSC was closely related to the LYSC sequence from other catfish species and located in the different clades with mammals and the goose-type lysozyme gene. The African catfish LYSC gDNA sequence was submitted to the Genbank under accession number: MN562879. Six SNPs were identified in African catfish LYSC. SNP1 was located in intron 1 (gDNA position 292, T/C), SNP2 and SNP3 were in intron 2 (position 889, T/C; and position 920, G/A), and SNP4, SNP5, and SNP6 were in intron 3 (position 1137, G/A; position 1237, G/T; and position 1280 G/C). LYSC polymorphism also found in other fish species. Japanese flounder *Paralichthys olivaceus* has eight polymorphism sites in its LYSC promoter and nine SNPs in the coding region (Liu et al. 2017). Asian seabass has two SNPs in the exon and the intron of the LYSC coding sequence (Fu et al. 2013).

### The mRNA expression and SNP associations to the *Aeromonas hydrophila* resistance

After the *A. hydrophila* challenge, the LYSC and other immune related-genes expression levels of resistant fish were higher than susceptible fish in several tissues ( $p < 0.05$ ), as shown in Figure 3. LYSC, LYSG, and HAMP expression were higher in the liver, head-kidney, and spleen ( $p < 0.05$ ), since the organs are reported to be the

major organs affected by *A. hydrophila* (Sahoo et al. 1998). The highest expression level of LYSC was observed in the liver, followed by head-kidney and spleen. The upregulation of LYSC and other immune genes expression after bacterial challenge also reported in channel catfish, Qihe crucian carp, and Darby's sturgeon after *A. hydrophila* challenge (Pridgeon et al. 2013; Wang et al. 2016; Zhang et al. 2018) and rohu carp after *Edwardsiella tarda* challenge (Mohanty dan Sahoo 2010) in the immune tissues. The expression of IL1B was higher in the head-kidney and spleen of susceptible fish (Figure 3;  $p < 0.05$ ). The excessive up-regulation of IL1B and other inflammatory cytokines showed an inefficient immune-response and lead to severe tissue damage, oxidative stress, and higher mortality in the susceptible group (Tort 2011; Dettleff et al. 2015). Upon *A. hydrophila* and other bacterial infections, the HAMP gene also showed significant expression, mainly in the liver of teleost fish (Das et al. 2015; Yang et al. 2007). This pattern might be due to the liver hepatocyte as the primary site for HAMP synthesis (Neves et al. 2016). These results suggest that the LYSC and HAMP gene might play an essential role in the fish effective immune responses against *A. hydrophila* infection in African catfish. Further research regarding the SNP in the HAMP gene is the potential to be done in the future.

Based on the LYSC genotyping results by sequencing among the seven resistant and five susceptible fish, we found three genotypes at SNP2: TT, CC, and CT (Figure 4). Susceptible fish were likely had more CC genotype than resistant fish and further analyzed with qPCR-HRM analysis to identify the genotype frequencies in SNP2. The results are shown in Table 3. In the resistant group, the frequencies of TT, CC, and TC were 27.78%, 5.55%, and 66.67%, respectively, while the frequencies of TT, CC, and TC were 16.67%, 27.78%, and 55.55% in the susceptible group, respectively. This result is suggesting that the SNP2 may be associated with the resistance to *A. hydrophila* infection in African catfish. In other fish species, the polymorphism in the LYSC gene was also reported to be associated with pathogenic bacteria resistance. Two SNPs in exon 2 (position 655, CC) and intron 3 (position 934, CT) of the LYSC gene reported being correlated with the big-belly disease resistance in Asian sea bass (Fu et al. 2013). The genotype CC at position 536, CC at position 1200, and GG at position 140 in the Japanese flounder LYSC were reported to be correlated with the resistance to *Listonela anguillarum* (Liu et al. 2017).

**Table 3.** The Distribution of SNP2 genotypes in resistant and susceptible groups of African catfish after *Aeromonas hydrophila* challenge

Population	Genotype	N total	Percentage	p-value
Resistant	TT	10	27.78	0.035
	CC	2	5.55	
	TC	24	66.67	
Susceptible	TT	6	16.67	0.035
	CC	10	27.78	
	TC	20	55.55	

**Start**

1 ATGAAGCCTT TGGTGTTCCT TCTGCTTCTG GCAGTTGTAA GTGCTAAAAA GTATGATCGG  
61 TGTGAGCTGG CAAGAGCCAT GAAAGCATTT GGCTTGGCTG GGTATCGTGG CATCAGTCTG  
121 GCTAACTgtg agtaaactgg tattttaaca tcctatgcag tacatctatc tgttgtgcaa  
181 aaggagctgg tgcagacaag gaagctgcag aaaattcagc ttgtttttat agctcaaagt  
241 gcactggtag ggtacatagg agtgacaaaa ctgataaaga taagataaaa tacttttta

**SNP1 c**

301 aatccagagg gaaactgtat atttaataata taatgcaaaa taagttaaata acataggaat  
361 aaatagtaca aatataaata tttcaataata taagaatatt taatttgcta aatatgtaca  
421 gtataagaaa taataaatta ttgattacca aattctgtct taccaggaca ataataataa  
481 taataatcaa atataacact gtacaatfff catactaata tacaataata gaactataac  
541 actataataa tatagaatff tacaccacaa aattacctaa caatatggtg tgtagtcaaa  
601 ttctgttttt tcccccgca aagtaactta aaaccttcta gaactaaaaa tccattttct  
661 ctgtttgttg tacagtgatt gtaataagta atgtatttct ataaatgctg aatattttct  
721 agGGGTTTGC TTGGCCAAAC ACGAGTTTGA CTTCAATACC AACGCCCAGA ACCGCAACTC  
781 TGATGGATCA ACAGACTATG GCATTTTCCA GATAACAAC CGTTGGTGGT GCACCAATGG  
841 GAAATTCCAT AGTGCTAACG GGTGCAAAC CTCCTGTAGT Ggtgagaat gttttttatt

**SNP2 c**

901 gatctattct tcaaatcttg aatgatgacc ataccaagct gtgaaaggct catatctttg

**SNP3 a**

961 tcttgatgcc tgcTTTTGCA aacagAGCTT CTTACCGATA ACATCGCGAA AGCTGTTGAG  
1021 TGTGCCAAGA CAATCGTGAG ACAGCAGGGA ATTACCGCAT Ggtaagacta ccttatacaat  
1081 atattgcatg tatggaggat gttgtaaaca tttagtgaag tctacacgtt tgagatcata

**SNP4 a**

1141 cttactgtat cataatcctt tagtttgcct tttttttcat tgcaaattac attaccggcc  
1201 ataactttta tgaaaagtaa aaatttcaga attgtggagt atttgttatt ttttggggg

**SNP5 t**


1261 ttattgacag tgtgcacttg agctgataat attcaatata aatttgctta tatttaaatt

**SNP6 c**


1321 gtaaacaaga aactgcaatg tctcacttat tgaatagtac agttgcacat tttctttatt  
1381 tgttgttgaa gtttaaaaat tgtaaaactt aatgttaatg atattaatgt gatcctaatt  
1441 ggactccaca gtagtgacgg tacatgattt attcattttc ctttttttac agGGTGGCAT  
1501 GCGTAACAA ATGCAGGGGT CGTGACGTAA GCTCTTACAC TGCAGGCTGT GGAGTTTAA

**Stop**


**Figure 1.** The gDNA sequence and the SNP position of African catfish LYSC gene. The capital letters are the exons, the small letters are the introns. Start and stop codons are underlined



**Figure 2.** The LYSC gene comparison between African catfish and other fish species. Exons are shown with boxes and introns with the line. The Genbank accession numbers are included under the species name



**Figure 3.** Relative mRNA expression levels of the LYSC and other immune-related genes of African catfish 24 h after *A. hydrophila* challenge. The  $\beta$ -actin gene was used for normalization and PBS control at 0 h as the expression calibrator. Data are shown as mean  $\pm$  SD (n= 4). Asterisks (\*) indicate the significant differences between resistant and susceptible fish (p<0.005).



**Figure 4.** Genotype variation and normalized melting plots of HRM analysis at SNP2 on the resistant and susceptible African fish after the *Aeromonas hydrophila* challenge. Three different genotypes were identified within the population: TT, CC, and TC.

For future development, it is essential to examine and confirm the association of the SNP2 in a larger population before its application as the potential marker in the African catfish selective breeding program. Moreover, the SNP2 association with other bacterial pathogens or stress resistance and its correlation with the growth traits could also be examined. In conclusion, the LYSC gDNA sequence was identified and characterized in African catfish. The modulation of its expression in the immune-tissues after *A. hydrophila* challenge suggests that the LYSC gene plays an essential role in *A. hydrophila* defense and resistance. The SNP2 found in the LYSC predicted to be associated with the resistance to *A. hydrophila* infection and may facilitate the future marker-assisted selection of the disease-resistance African catfish.


**ACKNOWLEDGMENTS**

There is no conflict of interest in this article. This work was supported by the Ministry of Research and Higher Education, Republic of Indonesia [grant number: PMDSU 1098/IT3.11/LT/2017] for Hasan Nasrullah and Alimuddin Alimuddin. We thank Dodi Hermawan and Dedi Supriyadi for their technical assistance in this study.

**REFERENCES**

Angka SL, Lam TJ, Sin YM. 1995. Some virulence characteristics of *Aeromonas hydrophila* in walking catfish (*Clarias gariepinus*). *Aquaculture*. 130 (2-3): 103-112. DOI: 10.1016/0044-8486(94)00216-B.

- Buonocore F, Randelli E, Trisolino P, Facchiano A, de Pascale D, Scapigliati G. 2014. Molecular characterization, gene structure, and antibacterial activity of a g-type lysozyme from the European sea bass (*Dicentrarchus labrax* L.). *Mol Immunol* 62 (1): 10-18. DOI: 10.1016/j.molimm.2014.05.009.
- Callewaert L, Michiels CW. 2010. Lysozymes in the animal kingdom. *J Biosci* 35 (1): 127-160. DOI: 10.1007/s12038-010-0015-5.
- Das A, Mohapatra A, Sahoo Pk. 2015. Cloning and characterization of antimicrobial peptide, hepcidin in Medium carp, *Puntius sarana*. *Int J Pept Res Ther* 21: 139-147. DOI: 10.1007/s10989-014-9438-4
- Dauda AB, Natrah I, Karim M, Kamarudin MS, Bichi A u H. 2018. African Catfish Aquaculture in Malaysia and Nigeria : Status, Trends, and African Catfish Aquaculture in Malaysia and Nigeria : Status, Trends, and Prospects. *Fish Aquacult J* 9 (1): 1-5. DOI: 10.4172/2150-3508.1000237.
- Detleff P, Bravo C, Patel A, Martinez V. 2015. Patterns of *Piscirickettsia salmonis* load in susceptible and resistant families of *Salmo salar*. *Fish Shellfish Immunol* 45 (1): 67-71. DOI: 10.1016/j.fsi.2015.03.039.
- Ekasari J, Suprayudi MA, Wiyoto W, Hazanah RF, Lenggara GS, Sulistiani R, Alkahfi M, Zairin M. 2016. Biofloc technology application in African catfish fingerling production: The effects on the reproductive performance of broodstock and the quality of eggs and larvae. *Aquaculture* 464: 349-356. DOI: 10.1016/j.aquaculture.2016.07.013.
- Di Falco F, Cammarata M, Vizzini A. 2017. Molecular characterisation, evolution and expression analysis of g-type lysozymes in *Ciona intestinalis*. *Dev Comp Immunol* 67: 457-463. DOI: 10.1016/j.dci.2016.09.010.
- Fu GH, Bai ZY, Xia JH, Liu F, Liu P, Yue GH. 2013. Analysis of Two Lysozyme Genes and Antimicrobial Functions of Their Recombinant Proteins in Asian Seabass. *PLoS One* 8 (11): e79743. DOI: 10.1371/journal.pone.0079743.
- Hanson LA, Liles MR, Hossain MJ, Griffin MJ, William G. 2014. Motile *Aeromonas* Septicemia. In: *American Fisheries Society-Fish Health Section. Blue Book*. Bethesda, Maryland.
- Hikima J, Hirono I, Aoki T. 2000. Molecular cloning and novel repeated sequences of a C-type Lysozyme Gene in Japanese Flounder (*Paralichthys olivaceus*). : 241-247. DOI: 10.1007/s101269900028.
- Liu J, Zhou N, Fu R, Cao D, Si Y, Li A, Zhao H, Zhang Q, Yu H. 2017. The polymorphism of the chicken-type lysozyme gene in Japanese flounder (*Paralichthys olivaceus*) and its association with resistance/susceptibility to *Listonella anguillarum*. *Fish Shellfish Immunol* 66: 43-49. DOI: 10.1016/j.fsi.2017.05.010.
- Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression data using real-time quantitative PCR and the 2- $\Delta\Delta$ CT method. *Methods* 25 (4): 402-408. DOI: 10.1006/meth.2001.1262.
- Mohanty BR, Sahoo PK. 2010. Immune responses and expression profiles of some immune-related genes in Indian major carp, *Labeo rohita* to *Edwardsiella tarda* infection. *Fish Shellfish Immunol* 28 (4): 613-621. DOI: 10.1016/j.fsi.2009.12.025.
- Nasrullah H, Nababan YI, Yanti DH, Hadiantho D, Nuryati S, Zairin Jr M, Eksari J, Alimuddin A. 2019. Identification and expression analysis of C-type and g-type lysozymes genes after *Aeromonas hydrophila* infection in African catfish. *J Akuakultur Indonesia* 18 (2): 1-10. DOI: 10.19027/jai.18.2.1-10.
- Neves, J, Caldas, C, Ramos, MF, Rodrigues, PNS. 2016. Hepcidin-Dependent Regulation of Erythropoiesis during Anemia in a Teleost Fish, *Dicentrarchus labrax*. *PLoS One*. 11: 1-17. DOI: 10.1371/journal.pone.0153940
- Pridgeon JW, Klesius PH, Dominowski PJ, Yancey RJ, Kievit MS. 2013. Chicken-type lysozyme in channel catfish: Expression analysis, lysozyme activity, and efficacy as immunostimulant against *Aeromonas hydrophila* infection. *Fish Shellfish Immunol* 35 (3): 680-688. DOI: 10.1016/j.fsi.2013.05.018.
- Sahoo PK, Mukherjee SC, Sahoo SK. 1998. *Aeromonas hydrophila* versus *Edwardsiella tarda*: a pathoanatomical study in *Clarias batrachus*. *J Aqua*. 6: 57-66
- Saurabh S, Sahoo PK. 2008. Lysozyme: An important defense molecule of fish innate immune system. *Aquacult Res* 39 (3): 223-239. DOI: 10.1111/j.1365-2109.2007.01883.x.
- Shoko AP, Limbu SM, Mgaya YD. 2016. Effect of stocking density on growth performance, survival, production, and financial benefits of African sharptooth catfish (*Clarias gariepinus*) monoculture in earthen ponds. *J Appl Aquacult* 28 (3): 220-234. DOI: 10.1080/10454438.2016.1188338.
- Tort L. 2011. Stress and immune modulation in fish. *Dev Comp Immunol* 35 (12): 1366-1375. DOI: 10.1016/j.dci.2011.07.002.
- Wang M, Zhao X, Kong X, Wang L, Jiao D, Zhang H. 2016. Molecular characterization and expressing analysis of the C-type and g-type lysozymes in Qihe crucian carp *Carassius auratus*. *Fish Shellfish Immunol* 52: 210-220. DOI: 10.1016/j.fsi.2016.03.040.
- Wang R, Feng J, Li C, Liu S, Zhang Y, Liu Z. 2013. Four lysozymes (one C-type and three g-type) in catfish are drastically but differentially induced after bacterial infection. *Fish Shellfish Immunol* 35 (1): 136-145. DOI: 10.1016/j.fsi.2013.04.014.
- Wei S, Huang Y, Huang X, Cai J, Wei J, Li P, Ouyang Z, Qin Q. 2014. Molecular cloning and characterization of a new G-type lysozyme gene (Ec-lysG) in orange-spotted grouper, *Epinephelus coioides*. *Dev Comp Immunol* 46 (2): 401-412. DOI: 10.1016/j.dci.2014.05.006.
- Yang M, Wang K, Chen J, Du H, Li S. 2007. Genomic organization and tissue-specific expression analysis of hepcidin-like genes from black porgy (*Acanthopagrus schlegelii* B.). *Fish Shellfish Immunol* 23: 1060-1071
- Yarahmadi P, Miandare HK, Fayaz S, Caipang CMA. 2016. Increased stocking density causes changes in expression of selected stress-and immune-related genes, humoral innate immune parameters, and stress responses of rainbow trout (*Oncorhynchus mykiss*). *Fish Shellfish Immunol* 48: 43-53. DOI: 10.1016/j.fsi.2015.11.007.
- Zhang S, Xu Q, Boscari E, Du H, Qi Z, Li Y, Huang J, Di J, Yue H, Li C, et al. 2018. Characterization and expression analysis of g-and C-type lysozymes in Dabry's sturgeon (*Acipenser dabryanus*). *Fish Shellfish Immunol* 76: 260-265. DOI: 10.1016/j.fsi.2018.03.006.



**Figure S1.** Phylogenetic tree of LYSC gDNA sequence of African catfish and other species. The tree was constructed using the neighbor-joining method with 1000 x bootstraps in MEGA X software